

RY reporting by health

ge 1 of CDER

ACT 1944. Date of birth: 62.7 kgs 82 Activerse Event or Product Problem Activerse Event or Problem Activerse Event or Initial Economic Information of prevent permanent or problem A pharmacist reported that a patient began taking an unspecified (mo/day/yr) 7/6/99 3. Describe event or problem A pharmacist reported that a patient began taking an unspecified regimen of acetaminophen on an unspecified date for an unspecified inficiation. The primary reporter stated that the patient took approximately 4 tablets the day pring to admission. On 2-Feb-99 the patient presented to the emergency care conter with JAUNDICE. ELEVATED INTERNATIONAL HORMALIZED RATIO (5.06). The primary reporter ison stated that the patient received one grain of acetaminophen in the emergency care conter prior to diagnosis. The primary reporter ison stated that the patient received one grain of acetaminophen in the emergency care conter prior to diagnosis. The primary reporter ison stated that the patient received one grain of acetaminophen in the emergency care conter prior to diagnosis. The patient was A DARTTED TO THE HOSPITAL, where troatment with acetyloystense was initiated. The patients's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 doses. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Serum Creatinine 1.6 2. Dose determine 1.6 2. Dose determine 1.6 3. Date of event problem 1.6 4. Date of the primary problem problem 1.6 Alteries Nikola Alteries Nikola Alteries days prior to admission, nausea and vomiting for one day prior or	e (give labeled strength mfr/labeler,	
Activerse Event or Product Problem Death Lie-shearening: Heaptiblization - initial Tongenital anomaly Heaptiblization - prolonged Disability Lie-shearening: Heaptiblization - prolonged Disability Congenital anomaly Required intervention to prevent permanent impairment/damage (mo/day/yr) Death Lie-shearening: Heaptiblization - prolonged Disability Activerse Event or problem A plasmacist reported that a patient began taking an unspecified regimen of acetaminophen on an unspecified date for an unspecified indication. The primary reporter stated that the patient took approximately 4 tablets the day prior to admission. On 2-Feb-99 the patient presented to the emergency care conter with JAUNDICE, LEVATED INTERNATIONAL HORMALIZED RATIO (5.90). The primary reporter also stated that the patient received one grain of acetaminephen in the emergency care conter prior to diagnosis. The patient was ADMITTED TO THE HOSPITAL, where treatment with patient was BDMITTED TO THE HOSPITAL, where treatment with acetalycystems was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Secum Creatinine: 1.6 272: INR 5.99. AST > 3200. ALT > 1600, acetaminophen level 15 4; 2/3: lotal bilirubin 4.9, INR 4.9 E. 1. Na. 7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse, patient presented with right abdominal pain, no oral intaky for three days prior to admission, nausea and vomibing for one day prior	and the second second second second second	if known)
n confidence) 3. Adverse Event or Product Problem 3. Adverse Event andre Product Problem 3. Adverse Event andre Product Problem 4. Disability Death Congenital anomaly Elephatization - initial Product Problem 5. Elephatization - prolonged Product Problem 6. Lie-theoatening Required intervention to prevent permanent impainment/damage 6. Lot Product Problem 7. Diste of event (mo/day/yr) 7. Describe event or problem 8. Diste of event (mo/day/yr) 8. Describe event or problem 8. Diste of event (mo/day/yr) 8. Describe event or problem 8. Apharmacist reported that a patient began taking an unspecified regimen of acetaminophen on an unspecified date for an unspecified indication. The primary reporter stated that the patient took approximately 4 lablets the day prine to admission. On 2-Feb-99 the patient presented to the emergency care conter with JAUNDICE, ELEVATED INTERNATIONAL HORMALIZED RATIO (5.06). The primary reporter islass stated that the patient received one grain of acetaminophen in the emergency care conter prior to diagnosis. The patient was ADMITTED TO THE INSPITAL, where troatment with acetylcysteina was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 doses. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates 8. Serum Creatinine: 1.6 227: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: itotal bilirubin 4.9, INR 4.9 7. Other relevant history, including preexisting medical conditions Allergies: INKOA alcohol abuse, patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior or ne day prior	cetaminophen	
Adverse Event or Product Problem Adverse Event and Product Problem Control of the Event and Product Problem Congenital anomaly Life-threatening: Life-threatening: Congenital anomaly Required intervention to prevent permanent impainment/damage (mo/day/yr) Date of event (mo/day/yr) Describe event or problem A pharmacist reported that a patient began taking an unspecified regimen of acetaminophen on an unspecified data for an unspecified regimen of acetaminophen on an unspecified data for an unspecified regimen of acetaminophen on an unspecified data for an unspecified regimen of acetaminophen on an unspecified approximately 4 lablets the day prior to admission. On 2-Feb-99 the patient presented to the memogracy care conter with JAUNDICE, ELEVATED LIVER ENYZMES (AST>3200, ALT>1600) and an ELEVATED INTERNATIONAL HORMALIZED RATIO (5.96). The patient presented is the emergency care conter prior to diagnosis. The patient was ADMITTED TO THE (DSPITAL, where treatment with acetylcysteria was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Serum Creatinine: 1.6] 2/2: (NR 5.96, AST > 3200, ALT > 1600, acetaminophen level 15 4; 2/3: itotal bilirubin 4.9, INR 4.9 Fe. 1. Na 7. Other relevant history, including preexisting medical conditions Allergies: (NKOA) alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior or need and prior or need to admission, nausea and vomiting for one day prior or need and prior or need to admission, nausea and vomiting for one day prior or need to a	- · · - · · · · · · · · · · · · · ·	
Adverse Event or Product Problem Adverse Event Image: Image	, frequency, route used	3. Therapy Dates (from/to)
Dutcennes attributed to adverse crisit Disability Product Problem #2 Dutcennes attributed to adverse crisit Disability Congenital anomaly #1 Lifetherateurg: Required intervention to prevent permanent impairment/damage #6 Date of event 2/3/99 4 Date of this report 7/6/99 #2 Date of event 2/3/99 4 Date of this report 7/6/99 #2 Describe event or problem Required intervention to prevent permanent impairment/damage #6 Date of event 2/3/99 4 Date of this report 7/6/99 #2 Describe event or problem Required for a unspecified #2 Describe event or problem Required date for an unspecified #6 Ediman of acetaminophen on an unspecified date for an unspecified egimen of acetaminophen on an unspecified date for an unspecified egimen of acetaminophen on an unspecified date for an unspecified egimen of acetaminophen on an unspecified date for an unspecified egimen of acetaminophen on an unspecified date for an unspecified egimen of acetaminophen on an unspecified date for an unspecified egimen victorial date egimen of acetaminophen on an unspecified date for an unspecified egimen victorial date egimen of acetaminophen on an unspecified egimen victorial date egimen victor		, _F ,
Disability Late-threatening: Life-threatening: L		
Death Life-threatening: Congenital anomaly Required intervention to prevent permanent impainment/damage Life-threatening: Required intervention to prevent permanent per vent permanent per vent permanent permanent per vent permanent per vent permanent per vent permanent permanent permanent (mo/day/yr) Describe event or problem A pharmacist reported that a patient began taking as unspecified permanent p		<u> </u>
Life-threatening: Congenital intervention to prevent permanent impainment/damage 10 Co.	nosis for use (indication)	5 Event abated after use stopped or dose reduced
Respitalization - initial prevent permanent impairment/damage initial		#1 Yes
Blospitalization - prolonged impainment/dainage		#2 1-3
3. Date of event (mo/day/yr) 4. Date of this report (mo/day/yr) 7.0799 3. Describe event or problem A pharmacist reported that a patient began taking an unspecified regimen of acetaminophen on an unspecified date for an unspecified indication. The primary reporter stated that the patient took approximately 4 tablets the day prior to admission. On 2-Feb-99 the patient presented to the emergency care conter with JAUNDICE, ELEVATED INTERNATIONAL NORMALIZED RATIO (5.96). The primary reporter risos stated that the patient eventure one grain of acetaminophen in the emergency care center prior to diagnosis. The patient was ADAITTED TO THE HOSPITAL, where treatment with acetylcysteine was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Serum Creatimine: 1.6 2/2: INR 5.96, AST > 3200, ALT > 1600, acetaminophen level 15 4; 2/3: total bilirubin 4.9, INR 4.9 E. 1. Na 7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intakt for three days prior to admission, nausea and vomiting for one day prior	(if known) 7. Exp. date	8. Event reappeared after
#2 Date of event (mo/day/yr) #2 Date of this report (mo/day/yr) #2 Professional	#1	reintroduction
(mo/day/yr) 5. Describe event or problem A pharmacist reported that a patient began taking an unspecified regimen of acetaminophen on an unspecified date for an unspecified indication. The primary reporter stated that the patient took approximately 4 tablets the day prior to admission. On 2-Feb-99 the patient presented to the emergency care conter with JAUNDICE, ELEVATED LIVER ENYZMES (AST>-3200, ALT>-1600) and an ELEVATED LIVER ENYZMES (AST>-3200, ALT>-1600) and an ELEVATED LIVER ENYZMES (AST>-3200, ALT>-1600) and an exterior reporter riso stated that the patient received one grain of acetaminophen in the emergency care center prior to diagnosis. The primary reporter riso stated that the patient received one grain of acetaminophen in the emergency care center prior to diagnosis. The patient was ABARTTED TO THE HOSPITAL, where treatment with acetylcysteine was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Serum Creatinine: 1.6 202: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15 4; 2/3: total bilirubin 4.9, INR 4.9 E. 1. Na 7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	b2	#1 Unknown
5. Describe event or problem A pharmacist reported that a patient began taking an unspecified egimen of acetaminophen on an unspecified date for an unspecified indication. The primary reporter stated that the patient took approximately 4 tablets the day prior to admission. On 2-Feb-99 the patient presented to the emergency care conter with JAUNDICE, ELEVATED LIVER ENYZMES (AST>-3200, ALT>-1600) and an ELEVATED LIVER ENYZMES (AST>-3200, ALT>-1600) and an ELEVATED LIVER NATIONAL ITORMALIZED RATIO (5.90). The primary reporter riso stated that the patient received one grain of acetaminophen in the emergency care center prior to diagnosis. The patient was ADMITTED TO THE HOSPITAL, where treatment with acetylcysteine was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Serum Creatinine: 1.6 202: INR 5.96, AST >-3200, ALT >-1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: NKDA altohol abuse; patient presented with right abdominal pain, no oral intakt for three days prior to admission, nausea and vomiting for one day prior	# (for product problems only)	#2
A pharmacist reported that a patient began taking an unspecified egimen of acetaminophen on an unspecified date for an unspecified indication. The primary reporter stated that the patient took epproximately 4 tablets the day prior to admission. On 2-Feb-99 the patient presented to the emergency care conter with JAUNDICE. ELEVATED LIVER ENYZMES (AST>3200, ALT>1600) and an ELEVATED INTERNATIONAL HORMALIZED RATIO (5.96). The primary reporter stated that the patient received one gran of acetaminophen in the emergency care center prior to diagnosis. The patient was ADMITTED TO THE HOSPITAL, where troatment with acetylcysteriae was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm crafty every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Scrum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: NKDA Allergies: NKDA Allergies: NKDA		<u></u>
egimen of acetaminophen on an unspective date to all disspections, indication. The primary reporter stated that the patient took approximately 4 tablets the day prior to admission. On 2-Feb-99 the patient presented to the emergency care conter with JAUNDICE, ELEVATED LIVER ENYZMES (AST-3200, ALT-1600) and an ELEVATED INTERNATIONAL NORMALIZED RATIO (5.96). The poimary reporter riso stated that the polient received one grain of acetaminophen in the emergency care center prior to diagnosis. The patient was ADMITTED TO THE HOSPITAL, where troatment with acetycysterae was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Scrum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	ncomitant modical products	r
poproximately 4 lablets the day prior to admission. On 2-response hip proximately 4 lablets the day prior to admission. On 2-response hip presented to the emergency care conter with JAUNDICE, LEVATED LIVER ENYZMES (AST>3200, ALT>1600) and an ELEVATED INTERNATIONAL HORMALIZED RATIO (5.96). The primary reporter islos stated that the patient received one grain of acetaminophen in the emergency care center prior to diagnosis. The natient was ADMITTED TO THE HOSPITAL, where troatment with acetylcysteina was initiated. The patient's regimen was 7.0 gm x1, then 3.5 gm crally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Serum Creatinine: 1.0 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior		
patient presented to the emergency care conter with Advicts. LEVATED LIVER ENTYMES (AST-3200, ALT>1600) and an ELEVATED INTERNATIONAL NORMALIZED RATIO (5.96). The brimary reporter idso stated that the polient received one grain of acetaminophen in the emergency care center prior to diagnosis. The natient was ADMITTED TO THE HOSPITAL, where troatment with acetycysteriae was initiated. The polient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Scrum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKDA altergies: NKDA altergies: NKDA altergies: NKDA		
LEVATED LIVER ENYZMES (ASTSZUT, ALT PRODUCTION) LEVATED INTERNATIONAL INCRMALIZED RATIO (5.96). The skimary reporter idso stated that the patient received one grain of incetaminophen in the emergency care center prior to diagnosis. The natient was ADMITTED TO THE HOSPITAL, where troatment with natient was fulfilled. The patient's regimen was 7.0 gm x 1, then cetylcysteine was initiated. The patient's regimen was 7.0 gm x 1, then not provided to the patient of the reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Secrum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: lotal bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: NKOA lacohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior		<u>. </u>
D. Social principles of the strip patient received the grain state of that the patient received the grain state of that the patient received the grain state of the patient prior to diagnosis. The patient was ADMITTED TO THE HOSPITAL, where treatment with acetylcysteriae was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Scrum Creatinine: 1.6 2/2: INR 5.96, AST > 3200, ALT > 1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKDA altergies: NKDA altergies: NKDA for three days prior to admission, nausea and vomiting for one day prior		
patient was ADMITTED TO THE HOSPITAL, where troatment with acetylcysteine was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 dosos. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Serum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15 4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: INKDA altochol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior		
patient was ADMRTTED THE HOSPITAL, when a 7.0 gm x 1, then accept/cysterine was initiated. The patient's regimen was 7.0 gm x 1, then 3.5 gm orally every 4 hours x 17 doses. The reaction was reported to have resolved. 6. Relevant tests/laboratory data, including dates Serum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15 4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: NKDA altechol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	136 Wal Davis	L
6. Relevant tests/laboratory data, including dates Serum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15 4; 2/3: lotal bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	uspect Medical Device	
6. Relevant tests/laboratory data, including dates Serum Creatinine 1.6 2/2: INR 5.96, AST >3200, ALT >1500, acataminophen level 15.4; 2/3: Itotal bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	These fields not used for electron	ic 3500 reporting at
6. Relevant tests/laboratory data, including dates Scrum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKOA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior		
6. Relevant tests/laboratory data, including dates Serum Creatinine 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acataminophen level 15 4; 2/3: total bilirubin 4.9, INR 4.9 E. 1. Na 7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intakt for three days prior to admission, nausea and vomiting for one day prior		
6. Relevant tests/laboratory data, including dates Scrum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	nternal ADR Event Co	ding
Serum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior		
Serum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: lotal bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	Reaction 1: increased in	
Serum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: lotal bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	```````	temational normalized ratio
Serum Creatinine: 1.6 2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: lotal bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	Reaction 3: jaundice	
2/2: INR 5.96, AST >3200, ALT >1600, acetaminophen level 15.4; 2/3: total bilirubin 4.9, INR 4.9 E. 7. Other relevant history, including preexisting medical conditions Altergies: NKOA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	Reaction 4:	
7. Other relevant history, including preexisting medical conditions Allergies: NKOA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	Reaction 5:	
7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	· -	
7. Other relevant history, including preexisting medical conditions Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	Reporter (see confidentia	ality section on bac
7. Other relevant history, including preexisting medical conditions Altergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	e, address and phone #	
Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	ADR Program Coordinator / Drug In	formation Service
Allergies: NKDA alcohol abuse; patient presented with right abdominal pain, no oral intake for three days prior to admission, nausea and vomiting for one day prior	Department of Pharmacy and Drug Is	
alcohol abuse; patient presented with right abdominal pain, no oral intakt for three days prior to admission, nausea and vomiting for one day prior		
for three days prior to admission, nausea and vomining for one day prior		
to adulation		
no admission 1 1 12 He	Ith Professional 3. Occupa	
	Yes [] No Pharmac	isl manufacturer
	nu do NOT want your identity	user facility
Mail to: MEDWATCH or FAX to:		distributor

JUL 2 0 1999

REC'D.

JUL 2 0 1999

MEDWATCH CTU